

University of Groningen

Optimal blockade of the renin angiotensin system in cardiorenal dysfunction

Wal, Rudy Martinus Adrianus van de

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2006

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Wal, R. M. A. V. D. (2006). *Optimal blockade of the renin angiotensin system in cardiorenal dysfunction*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

CHAPTER 1

General introduction Outline of the thesis

Ruud M.A. van de Wal
Adriaan A. Voors
Ron T. Gansevoort

Department of Cardiology, St. Antonius Hospital, Nieuwegein, The Netherlands
Department of Cardiology, University Medical Center Groningen, Groningen, The Netherlands
Department of Nephrology, University Medical Center Groningen, Groningen, The Netherlands

Urinary albumin excretion

Interest in the value of low levels of urinary albumin excretion (UAE) dates back to 1981, when it was shown that concentrations of urinary albumin that could not be detected by the standard dipsticks at that moment predict development of overt proteinuria in patients with diabetes mellitus.^{1,2} It was described that in such patients a urinary albumin excretion (UAE) between 30 and 140 mg/min had a 24 fold higher risk to develop nephropathy. A few years later, in 1984, it was also shown that such low levels of albuminuria predict mortality in patients with diabetes.³ Since, the term “microalbuminuria” has been introduced, reflecting the interest that rose in the clinical value of such slightly elevated levels of UAE.

In adults the normal mean value for UAE is 10 mg per day.⁴ However, UAE can be slightly increased under certain physiological circumstances, such as upright posture, exercise, pregnancy, and fever. Therefore, it is generally recommended in case an abnormal test result is obtained to confirm this on two separate occasions. The present definitions of microalbuminuria were described by the seventh report of the Joint National Committee (JNC 7) in 2002 and summarized in Table 1.

	Urine collection method	Normal	microalbuminuria	macroalbuminuria
Albumin	24-hour excretion	<30 mg/day	30-300 mg/day	>300 mg/day
	Spot urine albumin-Specific Dipstick	<20 mg/L	>20-200 mg/L	>200 mg/L
	Spot urine albumin-to creatinine ratio	<17 mg/g (♂) <25 mg/g (♀)	17-250 mg/g (♂) 25-355 mg/g (♀)	>250 mg/g (♂) >355 mg/g (♀)

Table 1. Definitions of albuminuria according to JNC 7 [4]

The glomerular capillary wall consists of fenestrated endothelium, the glomerular basement membrane, and the interdigitated foot processes of podocytes expressing the transmembrane protein nephrin.⁵ Under physiological conditions, the structural integrity of this filtration barrier prevents the abnormal passage of albumin (molecular mass 66 kDa) and high-molecular-weight proteins (>66 kDa), whereas low-molecular-weight proteins (<66 kDa) can pass without restriction. In addition, the transglomerular passage of macromolecules is regulated by the charge-selective properties of the

glomerular capillary membrane and the hemodynamic forces operating across the capillary wall.^{6;7}

In a healthy subject the amount of albumin excreted with urine normally represents less than 1% of the albumin filtered at the glomerular level.⁸ The remaining albumin is reabsorbed predominantly by the proximal tubuli through cellular mechanisms, by means of the synergistic receptors megalin and cubilin.⁹ This mechanism usually works quite close to saturation. As a consequence, any further increase in the amount of albumin filtered at the glomerular level will inevitably be accompanied by an increase in UAE.

In subjects with renal disease proteinuria is a good marker for renal disease progression.^{10;11} However, in renal disease urinary protein loss usually results from a specific lesion within the kidney, not per se reflecting the “health” status of the overall vasculature. Therefore, it remains unclear whether in these patients UAE can be utilized as a predictor of cardiovascular risk.

In hypertensive and diabetic patients albumin leakage is more often considered to be a reflection of generalized endothelial or vascular dysfunction,^{12;13} which is supported by the observation that elevated UAE is associated with elevated levels of high sensitivity C-reactive protein, von Willebrand factor and with impaired arterial dilatory capacity.¹⁴⁻¹⁶ It is hypothesized that damaged endothelium may lead to increased leakage of plasma albumin and lipids through the vessel walls and evoke an inflammatory response, thus linking elevated UAE to atherosclerosis.¹⁷ Since this process is not restricted to the kidney, microalbuminuria may be an indication of widespread vascular leakage of albumin. In addition to this hypothesis, increased intraglomerular pressure can also cause local renal leakage of albumin.

The evidence that albumin leakage is associated with cardiovascular risk is overwhelming. In recent studies on the potential prognostic value of microalbuminuria for cardiovascular events, the threshold value indicating increased risk has even been found to be well below the UAE values presently defining microalbuminuria regardless of the population included.¹⁸ A continuous cardiovascular risk spectrum has been observed starting from albumin excretion rates as low as 10 mg/24h.^{19;20}

Subsequent clinical evidence documented an association between UAE and other cardiovascular risk factors, target organ damage and risk of cardiovascular disease in the general population,²⁰⁻²³ and in specific clinical contexts, including essential hypertension.^{19;24;25} Accordingly, recent evidence suggests that reduction of UAE is associated with a lower cardiovascular risk.^{26;27}

Urinary albumin excretion and cardiovascular risk factors

Although UAE is a powerful predictor of cardiovascular morbidity and mortality, even today it continues to be unclear whether UAE is a risk factor per se, or merely a reflection of the effects of other risk factors (Figure 1). The fact is, that UAE is associated with biological factors like age and gender, but also often coincides with other risk factors, such as hypertension, insulin resistance, and increased levels of low-density lipoprotein.²⁸⁻³⁰

Hypertension

Over 30 years ago Parving et al. were the first to find an association between UAE and blood pressure in a hypertensive population.³¹ Since then several authors have confirmed this finding, and even in patients with high-normal blood pressure the prevalence of microalbuminuria has been shown to be increased.³² The prevalence of elevated UAE in hypertensive populations has been reported to be up to 46%, depending on the technique of measurement and the definition used.³³ Recent work suggests that hypertensive patients with increased UEA are more inclined to show early signs of other end-organ disease as well, including left ventricular hypertrophy and increased carotid artery thickness.³⁴ In addition, blood pressure lowering reduces UAE.




Figure 1. Possible factors involved in the pathogenesis of microalbuminuria (adapted from Verdecchia and Reboli).³³

The relation between insulin resistance and primary hypertension was described almost two decades ago.⁴⁶ Since then, several authors have investigated and discussed the association between insulin resistance and elevated UAE, but conflicting results were obtained.^{47;48} In addition, it remains uncertain whether their presumed association is independent of other factors, such as obesity and hypertension.

Lipids

In patients with hypertension and diabetes, the combined presence of elevated UAE and hyperlipidemia is frequent. In diabetic populations several authors reported an association between UAE and atherogenic lipids.⁴⁹⁻⁵¹ In two non-diabetic hypertensive populations microalbuminuric subjects had a significantly worse lipid profile than their non-albuminuric counterparts.^{52;53} In a large general population study, plasma cholesterol was an independent correlate of microalbuminuria.⁵⁴

Several explanations for the association between hyperlipidemia and elevated UAE have been proposed.⁵³ First, the rise in serum lipids could be caused, direct or indirect, by the loss of proteins that are involved in lipid production. Second, lipid abnormalities may contribute to glomerulosclerosis by a mechanism similar to atherosclerosis.⁵⁵ In support of this theory is the finding that lipid-lowering medication also ameliorates UAE,⁵⁶ although these results are not supported by others.²⁶

Smoking

Intrarenal hemodynamics can be influenced by smoking.⁵⁷ In smokers, the response of the kidney to increased systemic blood pressure may be impaired, possibly leading to increased intraglomerular capillary pressure.⁵⁸ The fact is, that smoking type I and type II diabetics excrete more albumin than non-smoking patients.^{59;60} In non-diabetics, hypertensive or not, smoking also seems to be independently associated with elevated UAE, even in the range defined as normal.⁶¹⁻⁶³ For this reason, other non-hemodynamic mechanisms must be involved in the pathophysiology of smoking-induced albumin leakage. Importantly, damage induced by smoking is probably not limited to renal endothelium, but throughout the body the endothelial function may be affected. Although several mechanisms have been proposed (e.g. alteration of prostaglandin/thromboxane pathway, generation of reactive oxygen species, carbon monoxide-induced hypoxia, tubulotoxicity, increased clotting of platelets, increased creatinin resistance), the exact underlying mechanism remains to be determined.⁶⁴

The underlying renal mechanism of hypertensive nephropathy has been the focus of research over the last years. At present, the common view is that due to failure of autoregulatory preglomerular constrictive response, the systemic hemodynamic load is conducted to the renal glomeruli.³⁵ The sympathetic nervous system may play a crucial role in this process.³⁶ Increased intraglomerular perfusion pressure will induce hyperfiltration and subsequently lead to a strain-related elevation of UAE.³⁷ Several studies have shown that agents that lower systolic and diastolic blood pressure reduce albuminuria, which establishes a strong argument in favor of this theory.³⁸⁻⁴¹ However, the finding that elevated UAE in hypertensive patients is an independent risk factor for cardiovascular disease, also suggests a link between vascular albumin leakage in the glomeruli, and systemic vascular damage. On the other hand, the pathogenesis of hypertensive nephropathy is multimodal and intrarenal mechanisms additional to blood pressure are involved, as the correlation with blood pressure can only partly explain urinary albumin loss. Interestingly, Brantsma et al recently reported that UAE may not always be merely the consequence of high blood pressure, but may also predict the development of hypertension.⁴²

Diabetes mellitus and insulin resistance

UAE, as a component of diabetic nephropathy, is a common complication in diabetic subjects with a prevalence between 10 to 42%. In the early stages the patient will show hyperfiltration, represented by a high glomerular filtration rate, and the occasional occurrence of microalbuminuria. Later, a gradual decline can be observed, while microalbuminuria progresses to macroalbuminuria.⁴³ Accordingly, an association between elevated UAE and disease duration has been reported.

Hyperglycemia induces renal cells to produce cytokines and growth factors that are thought to be responsible for structural changes and for functional changes, such as increased permeability of the glomerular basement membrane. Several humoral factors have been proposed to contribute to these modifications, such as transforming growth factor- β_1 (TGF- β_1), platelet-derived growth factor, connective tissue growth factor, protein kinase C, and vascular permeability factor.⁴⁴ Advanced glycosylation end products, which are abundantly present in the diabetic kidney, have also been recognized to induce extracellular matrix production and hence contribute to glomerular sclerosis.⁴⁵ According to the Steno hypothesis, this detrimental process is not restricted to renal tissue, but will, to a certain degree, also affect extrarenal matrix components.¹²

The renin angiotensin system

The renin angiotensin system (RAS) is an endocrine pathway which provides a homeostatic control mechanism for sodium balance, intravascular volume, and therefore blood pressure (Figure 2).^{65;66} The potent effector peptide, angiotensin II, is an octapeptide that is generated by cleavage from angiotensinogen through an action of two different peptides, renin and angiotensin converting enzyme (ACE).⁶⁷ The latter enzyme also links the RAS and the kallikrein-kinin system (KKS), as it breaks down bradykinin.⁶⁸




Figure 2. Simplified renin angiotensin system and targets for pharmacological intervention.

The role of angiotensin II in cardiovascular disease is well documented and nowadays this peptide is considered to be a key mediator of cardiovascular damage. The classical actions of angiotensin II include vasoconstriction, facilitation of sympathetic neurotransmission, and water and sodium retention. The latter can be accomplished directly or indirectly via aldosterone, a mineralocorticoid, which is considered to be an important mediator of angiotensin II-induced damage.⁶⁹

Angiotensin II mediates its effects through two main receptor subtypes: the angiotensin II type 1 receptor (AT1R) and the angiotensin II type 2 receptor (AT2R). These two receptors are expressed and distributed heterogeneously throughout the human body, and can be found in peripheral tissues, several organs (e.g. in the kidney) and the brain.⁷⁰ Both receptors are seven transmembrane

glycoproteins with only 32-34 % homology and they both have been cloned.^{71;72} The classical, in general detrimental, peripheral actions of angiotensin II are mediated by the AT1R (Table 2). Although knowledge is expanding rapidly, the (patho)physiological role of the AT2R is still poorly understood. Due to these effects angiotensin II contributes to the pathogenesis of vascular, cardiac, renal and cerebral pathologies, such as atherosclerosis, post-infarction remodeling, left ventricular hypertrophy, heart failure, stroke, and possibly diabetes.

AT1 receptor	AT2 receptor
Vasoconstriction	Vasodilatation (?)
Na ⁺ reabsorption/ H ₂ O retention	Apoptosis
Renin suppression	Tissue regeneration/repair
Inotropic effects	Inhibition of tissue proliferation
Vascular and cardiac hypertrophy (TGF- β_1)	Neuroprotection
Vascular injury and myocardial fibrosis	Cell differentiation
Proarrhythmic effects	Stimulation of bradykinin production
Prothrombotic effects (PAI-I \uparrow)	
Free radical formation (aging?)	
Facilitation of sympathetic transmission	
Endothelin secretion	
Proinflammatory effects	
Inhibition of Cell differentiation	
Facilitation of LDL transport	

Table 2. Angiotensin II mediated effects.
Abbreviations: TGF- β_1 , transformin growth factor β_1 ; PAI-1, plasminogen activator inhibitor 1

The traditional view of the RAS being a hormonal system, whereby angiotensin II is exclusively formed in the circulation and transferred to peripheral tissues, has long been rejected.^{73;74} A vast amount of evidence confirms that peripheral tissues are an important site of generation of this peptide, and local independent renin angiotensin systems have been found in many tissues, including brain, kidney, adrenal gland, pancreas, testis, blood vessels, and the heart.⁷⁰ Importantly, the autocrine and paracrine effects contribute to the regulation of local cellular and tissue functions, independent of the circulating endocrine system.⁷⁵ Furthermore, It is important to note that even within organs tissue RAS can be locally compartmentalized. For instance, within the kidney overall angiotensin II concentrations are higher in the medulla compared with the cortex.⁷⁶

Renal and vascular endothelial effects of the RAS

RAS and Kidney

Sodium retention and excretion, resulting from aldosterone which is released after activation of AT1R in the kidney, provokes change of blood pressure and controls volume status. Other effects mediated via AT1R include regulation of endocrine functions, and stimulation of mitogenic pathways.

Angiotensin II exerts specific actions on intrarenal hemodynamics, thus contributing to a higher perfusion pressure. Angiotensin II-induced constriction of the efferent arteriole reduces renal blood flow and raises glomerular capillary pressure, consequently augmenting glomerular filtration rate and filtration fraction.^{77;78} The changes in oncotic (\uparrow) and hydrostatic (\downarrow) pressure which are thus created in the peritubular vessels, are transduced to the interstitium, and promote a shift of sodium and water from the proximal tubule into the interstitium and systemic circulation. The reduction of medullary perfusion and diminished interstitial pressure simultaneously lower sodium and water excretion.⁷⁸ Beside these hemodynamic effects, the RAS increases water and sodium resorption by acting directly on several ion pumps (e.g. Na^+/H^+ -antiporter, Na^+/K^+ -ATPase, $\text{Na}^+/\text{HCO}_3^-$ -cotransporter) located throughout the tubular system.^{79;80} Thus, the RAS plays a pivotal role in maintaining normovolumic state and a normal ion balance. However, derangement of this hormonal cascade is thought to induce renal dysfunction (Figure 3).




Figure 3. Our knowledge of the relation between the local and circulating renin angiotensin system, arterial blood pressure and urinary albumin excretion remains incomplete.

Furthermore, angiotensin II is responsible for the production of nephrotoxic oxygen species, profibrotic cytokins, and growth factors, consequently inducing cell proliferation and tissue remodeling. Together these effects promote the development of glomerulosclerosis and tubulointerstitial fibrosis.^{81;82} Renal remodeling is augmented by angiotensin-induced aldosterone synthesis, as this hormone, besides being an important regulator of water and salt homeostasis, shares the ability to provoke mitogenic and profibrotic changes.⁸³ TGF- β_1 , plasminogen activator inhibitor-1, reactive oxygen species, endothelial dysfunction, upregulation of the AT1R have all been proposed as possible mediators of angiotensin II- and aldosterone-mediated renal injury and scarring (Table 2).⁸³ However, only few reports have been published showing a direct relationship between angiotensin II and glomerular permselectivity.^{84;85}

The above-mentioned effects may induce and maintain glomerular leakiness, and therefore play a role in UAE. Conversely, urinary albumin may also stimulate the RAS in proximal tubular cells.⁸⁶

Hemodynamic effect	Non-hemodynamic effect
Increased glomerular capillary pressure:	Induction of renal hypertrophy and cell proliferation
1. Post glomerular vasoconstriction (direct or indirect via endothelins \uparrow or NO \downarrow)	Modulation of extracellular matrix synthesis (\uparrow) and degradation (\downarrow)
2. Systemic hypertension	Stimulation of cytokine (e.g. TGF- β_1 , VEGF, PAI-1)
Filtration surface area reduction due to mesangial cell contraction	Production of free oxygen radicals

Table 3. Summary of proposed angiotensin II-mediated mechanisms leading to increased urinary albumin excretion (adapted from Leehey et al.⁸⁷).
Abbreviations: TGF- β_1 , transforming growth factor- β_1 ; VEGF, vascular endothelial growth factor; PAI-1, plasminogen activator inhibitor-1; NO; nitric oxide

RAS and vascular endothelium

Angiotensin II exerts its proinflammatory effects through the AT1R (Table 2).⁸⁸ One of the most important, angiotensin II-modulated steps, in the initial stage of inflammation is the increase in vascular permeability.⁸⁹ Increased permeability of the vascular endothelium probably results from pressure-mediated mechanical injury, but also from angiotensin II-induced second mediators that may influence permeability (VEGF, prostaglandins).⁸⁸ This important early manifestation of inflammation and atherosclerosis will subsequently lead to cell infiltration and

exudation of protein-rich fluid. In line with this theory, an association between plasma renin activity and UAE was described in a small study performed in a population with essential hypertension.⁹⁰ In addition, a few authors found an association between the angiotensin converting enzyme DD genotype and the presence of elevated UAE.⁹¹⁻⁹³ For other genetic variants within the RAS (AGT M235T, and AT1 A1266C gene polymorphisms) this relationship remains to be determined.^{93;94}

Pharmacological inhibition of the RAS in renal dysfunction

The cumulative incidence of diabetic nephropathy after duration of either type I or type II diabetes of 25 years has been reported to be 25-40%.⁹⁵ As diabetic and hypertensive nephropathy are the leading causes of end-stage renal disease in Europe and the United States, it is important to prevent, or at least to delay, in such subjects disease progression. Moreover, in non-diabetics the degree of UAE is related to cardiovascular prognosis. Considering this and the rising incidence of type II diabetes mellitus and its complications, it is imperative to develop new and to optimize old pharmacological strategies. Subsequently, the level of UAE may be used to monitor treatment efficacy,⁹⁶ although heterogeneity of changes in UAE during treatment may complicate the interpretation of efficacy.

ACE inhibitors

The antihypertensive effects of the first compound blocking the RAS were described in the early 1970s.^{97;98} A few years later the first orally available ACE inhibitor, captopril, was marketed. Nowadays, more than 10 different ACE inhibitors are available for human use, all having specific pharmacological properties. ACE inhibitors block circulating and tissue ACE, and therefore lower systemic and local angiotensin II production and raise bradykinin levels (Figure 1). Besides lowering blood pressure, ACE inhibitors have clearly demonstrated to reduce proteinuria in both diabetic and non-diabetic nephropathy.⁹⁹ Several meta-analyses, suggest that this antiproteinuric effect is partly independent of blood pressure reduction, and greater than can be achieved with calcium blocking agents, diuretics and beta-blockers.⁹⁹ This suggests additional RAS-related effect up-and-above hemodynamic effects.

The Diabetic Nephropathy Trialist Group published a meta-analysis in which they demonstrate that ACE inhibitors significantly slow down the progression from microalbuminuria to macroalbuminuria in (type I) diabetic patients.¹⁰⁰

In patients with overt nephropathy of type 1 diabetes, ACE inhibitors not only lowered UAE, but in addition significantly delayed the decline in glomerular filtration rate that is commonly observed in diabetic patients suffering from nephropathy.^{101;102}

Data from the diabetic population (primarily type II diabetes mellitus) of the Heart Outcomes Prevention Evaluation (HOPE) trial suggested the ACE inhibitor ramipril, compared with placebo, significantly reduced the risk of developing nephropathy in these subjects.¹⁰³ Recently, the Bergamo Nephrologic Diabetes Complications Trial (BENEDICT) indicated that in hypertensive, normo-albuminuric type II diabetics the ACE inhibitor trandolapril was even more effective in preventing the development of microalbuminuria than the non-dihydropyridine calcium channel blocker verapamil.¹⁰⁴ In contrast, a small study in hypertensive type II diabetics did not reveal a significant difference between the antialbuminuric effects of ramipril and lercanidipine.¹⁰⁵

ACE inhibitors have a clear renoprotective effect in non-diabetic renal disease, when compared to compounds not interfering with the activity of the RAS.¹⁰⁶ In patients with non-diabetic renal disease use of ramipril was even associated with an improvement of glomerular filtration rate.¹⁰⁷ The renoprotective effects of ACE inhibitors are probably mediated by their specific ability to block angiotensin II-induced constriction of the post-glomerular capillaries. The resulting fall in glomerular capillary pressure consequently reduces glomerular filtration rate and filtration fraction.

Beside their specific renoprotective effect, ACE inhibitors also protect the cardiovascular system. There is much debate whether ACE inhibitors offer more cardiovascular protection than the older blood pressure lowering agents, diuretics and beta-blockers. The meta-regression analysis by Staessen et al suggests that the efficacy of blood pressure lowering agents to prevent cardiovascular accidents and myocardial infarction is predominantly dependent on the reduction in blood pressure that is achieved.¹⁰⁸ Some recent individual studies suggests however, that ACE inhibitors may confer a cardiovascular protective effect that goes beyond blood pressure reduction per se.^{103;109-112} Interestingly, a number of post-hoc analyses attributed the superior CV effect of ACE inhibition to be dependent on the level of albuminuria, for instance in the HOPE study, in which patients at high risk for cardiovascular events were randomised to either ramipril or placebo. Subgroup analysis of this study showed that in those patients with high baseline UAE, intervention with an ACE inhibitor is of particular value and improves cardiovascular prognosis.¹⁰³ The potency of

ACE inhibitors to induce cardiovascular benefits is also illustrated by the results from the Prevention of Vascular and Endstage Renal Disease Intervention Trial (PREVEND-IT). In this trial performed in non-hypertensive microalbuminuric subjects, fosinopril reduced UAE significantly and in addition, treatment was associated with a trend in reducing cardiovascular events.²⁶ Interestingly, the efficacy of fosinopril in this study to prevent cardiovascular events was shown to be dependent on baseline UAE. In subjects with UAE>50 mg/24hr the ACEi induced a 60% relative risk reduction in CV endpoints, whereas in subjects with baseline UAE 15-50 mg/24hr almost no cardioprotective effect was noted.

By blocking angiotensin II generation, ACE inhibitors induce a postglomerular vasodilation as stated above. This will reduce intraglomerular pressure and may consequently lead to a reversible reduction of GFR. In turn, this will lead to renoprotection in the long run (“short term pain, is long term gain”). Unfortunately, this may force a small selection of patients to suspend or discontinue therapy. Hypotension, cough and hyperkalemia are relatively common side effects, whereas angioneurotic edema is a rare but potentially dangerous side effect of ACE inhibitors.

In summary, ACE inhibitors have demonstrated to be valuable antiproteinuric and renoprotective agents in both diabetic and non-diabetic subjects. Over the last few years evidence for their cardiovascular protective effects is mounting as well. Some evidence even suggests that these agents exert an extra beneficial cardiovascular profile in comparison to other blood pressure lowering agents, especially in albuminuric subjects.

Angiotensin II receptor blockers

The first peptide angiotensin type I receptor blockers (ARB), saralazin, was described in 1971,¹¹³ but it took another 20 years before the first non-peptide ARB, losartan, became available for patient use. Since then, several ARBs have proven to lower blood pressure, slow down the progression of diabetic and non-diabetic renal disease, reduce proteinuria irrespective of the type of renal disease, and also reduce the risk of overt nephropathy.^{114;115}

	Treatment	Patients	Follow up	Primary endpoint	Outcome
Parving HH, 2001	Irbesartan 150 mg 300 mg Placebo	590 patients, MA Hypertension	2y	Time to onset diabetic nephropathy	I150 vs P HR=0.56 I300 vs P HR=0.32
Lewis EJ, 2001	Irbesartan 200 mg Amlodipine 10 mg Placebo	1715 patients Proteinuria Hypertension	2.6y	Doubling serum creat Development of ESRD Death from any cause	I vs A RR=0.77 I vs P RR=0.80
Brenner MB, 2001	Losartan 50-100 mg Placebo	1513 patients Proteinuria	3.4y	Doubling serum creat Development of ESRD Death from any cause	L vs P RR=0.84
Viberti G, 2002	Valsartan 80 mg Amlodipine 5 mg	332 Patients, MA	1y	% Change UAE from baseline	V:56% UAE ↓ A: 8% UAE ↓

Table 4. Randomized controlled trials performed with angiotensin II receptor blockers in type II diabetic patients.

Abbreviations: MA; microalbuminuria, creat; creatinine, ESRD; end-stage renal disease, HR; hazard ratio, RR; relative risk

In contrast to ACE inhibitors, the evidence for a beneficial ARB-induced renoprotective effect in type 1 diabetic patients is limited.¹¹⁶ Although ACE inhibitors and ARBs seem to have comparable effects on renal hemodynamics,¹¹⁷ only small randomised placebo-controlled trials have actually studied the renoprotective properties of ARBs in type 1 diabetes.¹¹⁸

In type 2 diabetic patients, however, the evidence is compelling. Four large randomised clinical trials (IRMA-2,¹¹⁹ IDNT,¹²⁰ RENAAL,¹²¹ MARVAL¹²²) established the beneficial (dose-dependent) effects of ARBs on a variety of renal endpoints in early and more advanced diabetic nephropathy (Table 4). Data from several other studies support these results.¹¹⁵ Importantly, treatment with an ARB is not inferior to ACE inhibition, when renoprotective efficacies are compared.¹²³

The antiproteinuric effects of ARBs in early non-diabetic hypertensive nephropathy have been described in animal experiments.^{124;125} Initially, only smaller randomised clinical trials were performed in hypertensive non-diabetic subjects with¹²⁶ and without advanced renal disease.¹²⁷⁻¹³² Later, post-hoc analysis of a larger cohort of 918 patients with isolated systolic hypertension demonstrated that telmisartan reduced UAE irrespective of baseline UAE,

and to a greater extent than hydrochlorothiazide.¹³³ The most convincing evidence was provided by the Losartan Intervention For Endpoint Reduction in Hypertension (LIFE) study.¹³⁴ In 8602 hypertensive patients with ECG-ascertained left ventricular hypertrophy, treatment with losartan significantly reduced UAE. In addition, the reduction was significantly greater than the one achieved with atenolol, as was the reduction in cardiovascular endpoints. The authors concluded that part of the mechanism behind the superiority of losartan is related to its greater UAE-reducing effect.¹³⁴

Interestingly, ARBs seem to have pleiotropic effects. For example, telmisartan has the ability to interact with the nuclear peroxisome proliferator-activated receptor gamma (PPAR- γ),¹³⁵ and olmesartan has been reported to exhibit anti-inflammatory effects in hypertensive patients.¹³⁶ The clinical value of these findings needs further exploration.

In summary, both ACE inhibitors and ARBs are first choice components of an antihypertensive treatment regimen in patients with diabetic or non-diabetic renal disease with albuminuria.^{114;137} However, the predicted specific cardiovascular benefits of RAS-inhibitors in terms of survival and morbidity in subjects with renal damage need further verification.

Aldosterone receptor antagonists

The role of aldosterone in the development and progression of renal injury has been studied in variety of experimental models of progressive (hypertensive) renal disease.¹³⁸⁻¹⁴⁰ Rodent experiments suggest that local overproduction of the mineralocorticoid receptor may be a significant factor in the predisposition to hypertension and the subsequent vascular and renal injuries.¹⁴¹ In agreement with these findings, patients with an aldosterone-secreting tumour often have proteinuria.¹⁴² Furthermore, several studies provide evidence that spironolactone, a non-selective aldosterone receptor antagonist, has a renovascular protective effect.¹⁴³⁻¹⁴⁵ Spironolactone also improved left ventricular remodeling in patients experiencing a first anterior myocardial infarction,¹⁴⁶ and reduced cardiovascular mortality in severe chronic heart failure patients.¹⁴⁷ Recent data suggest that spironolactone may also improve acetylcholine induced vasodilation and thus protect endothelial function in patients with mild (NYHA class I or II) heart failure.¹⁴⁸ However, due to the lack of selectivity for the mineralocorticoid receptor and its endocrine side effects, the widespread use of spironolactone in humans is limited.

The selective aldosterone receptor antagonist eplerenone has recently demonstrated to have renoprotective properties that go beyond its antihypertensive effects as well. Compared with enalapril¹⁴⁹ and amlodipine,¹⁵⁰ the antiproteinuric properties of eplerenone were significantly better. In addition, combination of eplerenone and enalapril reduced UAE even to a greater extent than therapy with either drug alone.¹⁴⁹ However, whether this translates into long-term cardiovascular protection remains to be established. Although eplerenone has less antiandrogenic side effects than spironolactone, both drugs increase serum potassium, especially in patients using an ACE inhibitor or an ARB. Ensuing hyperkalemia, although troublesome, is reversible, and depends on dose and creatinine clearance.¹⁵¹ The fact that eplerenone is better tolerated than spironolactone offers the possibility to use eplerenone alone, or combined with an ACE inhibitor or ARB as a new renoprotective strategy. In the EPHESUS trial eplerenone demonstrated to improve post-infarction left ventricular remodeling, and subsequently reduce cardiovascular mortality.¹⁵² Initial reports, although performed in rodents, suggest that the cardioprotective properties of this agent may be mediated, in part, by stimulating endothelial NO synthase.¹⁵³ Extrapolation of these results to patients without heart failure is complex, and currently unfeasible. However, these data again indicate that agents that interfere in the renin-angiotensin-system may result in reno- and cardiovascular protection that goes beyond what might be expected from blood pressure lowering alone.

Renin inhibitors

The latest addition to the RAS-blocking armamentarium are the renin inhibitors.¹⁵⁴ After a few disappointing predecessors, Aliskiren (SPP100) seems to be the first potent orally active alkane carboxamide renin inhibitor and the first reports suggest that its blood pressure lowering effects are comparable to other RAS blocking agents.^{155;156} In addition, aliskiren is well-tolerated.¹⁵⁵ By actively antagonizing the rate-limiting step of the RAS cascade a general down-regulation of RAS activity can be observed.¹⁵⁶⁻¹⁵⁸ Interestingly, Pilz et al. demonstrated that both low and high dose aliskiren reversed albuminuria and normalized serum creatinine in transgenic rats.¹⁵⁹ New renin inhibitors may therefore become an alternative to ACE inhibitors and ARBs in the treatment of hypertension, and possibly also provide end organ protection. Other renin inhibitors, for example zalkiren,¹⁶⁰ are currently under investigation. To date, none of these agents has proven to be cardioprotective.

Conclusion

Increased UAE is a common finding in diabetic patients, as well as in hypertensive patients without renal disease, which negatively affects prognosis. Even in the general population subjects with increased UAE experience more cardiovascular events than subjects with normal UAE. Although increased UAE seems to be interrelated with several other cardiovascular risk factors and markers, evidence emerges that albuminuria is also an independent risk factor. As blocking the RAS induces a specific reduction of UAE that is greater than might be expected from blood pressure lowering alone, a causal relationship between RAS and UAE is suggested. Beside ACE inhibitors and blockers of the AT1R, (selective) aldosterone receptor antagonists and renin inhibitors may also be able to reduce UAE. Yet, whether the efficacy of these compounds to lower albuminuria is a specific renal mechanism, or a reflection of a superior efficacy of these drugs to ameliorate generalized endothelial dysfunction is yet unknown. Recent evidence suggests however, that these compounds may have a specific cardioprotective effect in subjects with higher levels of albuminuria, besides their proven renoprotective value.

Outline of this thesis

This present thesis concerns dysfunction of the cardiorenal axis, with a special focus on elevated urinary albumin excretion and the renin angiotensin system. The effect of pharmacological blockade of the renin angiotensin system on the activity of the renin angiotensin system and on urinary albumin excretion will also be discussed.

The value of pre-operative renal function as a predictor of long-term outcome in patients undergoing coronary artery bypass graft surgery will be described in **Chapter 2**. In **Chapter 3** we describe the prevalence of microalbuminuria in a cohort of severe chronic heart failure patients. We also present the neurohormonal variation that coincides with elevated urinary albumin excretion in these patients. The renin angiotensin system can be blocked by ACE inhibitors and angiotensin II type 1 receptor blockers. **Chapter 4** elaborates on the renin angiotensin system in chronic heart failure patients and the effect of ACE inhibition on the activity of this cascade. One of the benefits attributed to angiotensin II type 1 receptor blockers is parallel stimulation of the angiotensin II type 2 receptor. For this reason, we analyse the vasoactive effects of the type 2 receptor in human arteries in **Chapter 5**. Several studies suggest that combination of angiotensin II type 1 receptor blockers and ACE inhibitors can provide a more complete protection against the detrimental effects of the renin angiotensin system. In **Chapter 6** we discuss the pros and cons of this strategy. Finally, in **Chapter 7** we present data from the PREVEND intervention trial. In otherwise healthy, microalbuminuric subjects we determine which parameters affect baseline urinary albumin excretion. In an additional analysis we verify the antialbuminuric effect of ACE inhibition, and we describe which parameters determine the efficacy of treatment.

References

1. Viberti GC, Hill RD, Jarrett RJ, Argyropoulos A, Mahmud U, Keen H: Microalbuminuria as a predictor of clinical nephropathy in insulin-dependent diabetes mellitus. *Lancet* 1982, 1:1430-1432.
2. Svendsen PA, Oxenboll B, Christiansen JS: Microalbuminuria in diabetic patients--a longitudinal study. *Acta Endocrinol.Suppl (Copenh)* 1981, 242:53-54.
3. Mogensen CE: Microalbuminuria predicts clinical proteinuria and early mortality in maturity-onset diabetes. *N.Engl.J.Med.* 1984, 310:356-360.
4. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am.J.Kidney Dis.* 2002, 39:S1-266.
5. Deen WM: What determines glomerular capillary permeability? *J.Clin.Invest* 2004, 114:1412-1414.
6. Brenner BM, Hostetter TH, Humes HD: Molecular basis of proteinuria of glomerular origin. *N.Engl.J.Med.* 1978, 298:826-833.
7. Brenner BM, Bohrer MP, Baylis C, Deen WM: Determinants of glomerular permselectivity: Insights derived from observations in vivo. *Kidney Int.* 1977, 12:229-237.
8. Rowe DJ, Dawnay A, Watts GF: Microalbuminuria in diabetes mellitus: review and recommendations for the measurement of albumin in urine. *Ann.Clin.Biochem.* 1990, 27 (Pt 4):297-312.
9. Christensen EI, Birn H: Megalin and cubilin: synergistic endocytic receptors in renal proximal tubule. *Am.J.Physiol Renal Physiol* 2001, 280:F562-F573.
10. Iseki K, Ikemiya Y, Iseki C, Takishita S: Proteinuria and the risk of developing end-stage renal disease. *Kidney Int.* 2003, 63:1468-1474.
11. Verhave JC, Gansevoort RT, Hillege HL, Bakker SJ, de Zeeuw D, de Jong PE: An elevated urinary albumin excretion predicts de novo development of renal function impairment in the general population. *Kidney Int.Suppl* 2004,S18-S21.
12. Deckert T, Feldt-Rasmussen B, Borch-Johnsen K, Jensen T, Kofoed-Enevoldsen A: Albuminuria reflects widespread vascular damage. The Steno hypothesis. *Diabetologia* 1989, 32:219-226.
13. Jensen JS: Renal and systemic transvascular albumin leakage in severe atherosclerosis. *Arterioscler.Thromb.Vasc.Biol.* 1995, 15:1324-1329.
14. Festa A, D'Agostino R, Howard G, Mykkanen L, Tracy RP, Haffner SM: Inflammation and microalbuminuria in nondiabetic and type 2 diabetic subjects: The Insulin Resistance Atherosclerosis Study. *Kidney Int.* 2000, 58:1703-1710.
15. Stehouwer CD, Nauta JJ, Zeldenrust GC, Hackeng WH, Donker AJ, den Ottolander GJ: Urinary albumin excretion, cardiovascular disease, and endothelial dysfunction in non-insulin-dependent diabetes mellitus. *Lancet* 1992, 340:319-323.
16. Clausen P, Jensen JS, Jensen G, Borch-Johnsen K, Feldt-Rasmussen B: Elevated urinary albumin excretion is associated with impaired arterial dilatory capacity in clinically healthy subjects. *Circulation* 2001, 103:1869-1874.
17. Jensen JS, Borch-Johnsen K, Jensen G, Feldt-Rasmussen B: Microalbuminuria reflects a generalized transvascular albumin leakiness in clinically healthy subjects. *Clin.Sci.(Lond)* 1995, 88:629-633.
18. Klausen KP, Scharling H, Jensen G, Jensen JS: New definition of microalbuminuria in hypertensive subjects: association with incident coronary heart disease and death. *Hypertension* 2005, 46:33-37.

19. Gerstein HC, Mann JF, Yi Q, Zinman B, Dinneen SF, Hoogwerf B, Halle JP, Young J, Rashkow A, Joyce C, Nawaz S, Yusuf S: Albuminuria and risk of cardiovascular events, death, and heart failure in diabetic and nondiabetic individuals. *JAMA* 2001, 286:421-426.
20. Hillege HL, Fidler V, Diercks GF, van Gilst WH, de Zeeuw D, van Veldhuisen DJ, Gans RO, Janssen WM, Grobbee DE, de Jong PE: Urinary albumin excretion predicts cardiovascular and noncardiovascular mortality in general population. *Circulation* 2002, 106:1777-1782.
21. Borch-Johnsen K, Feldt-Rasmussen B, Strandgaard S, Schroll M, Jensen JS: Urinary albumin excretion. An independent predictor of ischemic heart disease. *Arterioscler.Thromb.Vasc.Biol.* 1999, 19:1992-1997.
22. Romundstad S, Holmen J, Kvenild K, Hallan H, Ellekjaer H: Microalbuminuria and all-cause mortality in 2,089 apparently healthy individuals: A 4.4-year follow-up study. The Nord-Trondelag Health Study (HUNT), Norway. *Am.J.Kidney Dis.* 2003, 42:466-473.
23. Damsgaard EM, Froland A, Jorgensen OD, Mogensen CE: Microalbuminuria as predictor of increased mortality in elderly people. *BMJ* 1990, 300:297-300.
24. Jensen JS, Feldt-Rasmussen B, Strandgaard S, Schroll M, Borch-Johnsen K: Arterial hypertension, microalbuminuria, and risk of ischemic heart disease. *Hypertension* 2000, 35:898-903.
25. Wachtell K, Ibsen H, Olsen MH, Borch-Johnsen K, Lindholm LH, Mogensen CE, Dahlof B, Devereux RB, Beevers G, de Faire U, Fyhrquist F, Julius S, Kjeldsen SE, Kristianson K, Lederballe-Pedersen O, Nieminen MS, Okin PM, Omvik P, Oparil S, Wedel H, Snapinn SM, Aurup P: Albuminuria and cardiovascular risk in hypertensive patients with left ventricular hypertrophy: the LIFE study. *Ann.Intern.Med.* 2003, 139:901-906.
26. Asselbergs FW, Diercks GF, Hillege HL, van Boven AJ, Janssen WM, Voors AA, de Zeeuw D, de Jong PE, van Veldhuisen DJ, van Gilst WH: Effects of fosinopril and pravastatin on cardiovascular events in subjects with microalbuminuria. *Circulation* 2004, 110:2809-2816.
27. Ibsen H, Olsen MH, Wachtell K, Borch-Johnsen K, Lindholm LH, Mogensen CE, Dahlof B, Devereux RB, de Faire U, Fyhrquist F, Julius S, Kjeldsen SE, Lederballe-Pedersen O, Nieminen MS, Omvik P, Oparil S, Wan Y: Reduction in albuminuria translates to reduction in cardiovascular events in hypertensive patients: losartan intervention for endpoint reduction in hypertension study. *Hypertension* 2005, 45:198-202.
28. Verhave JC, Hillege HL, Burgerhof JG, Navis G, de Zeeuw D, de Jong PE: Cardiovascular risk factors are differently associated with urinary albumin excretion in men and women. *J.Am.Soc.Nephrol.* 2003, 14:1330-1335.
29. Rosa TT, Palatini P: Clinical value of microalbuminuria in hypertension. *J.Hypertens.* 2000, 18:645-654.
30. Jensen JS, Feldt-Rasmussen B, Borch-Johnsen K, Clausen P, Appleyard M, Jensen G: Microalbuminuria and its relation to cardiovascular disease and risk factors. A population-based study of 1254 hypertensive individuals. *J.Hum.Hypertens.* 1997, 11:727-732.
31. Parving HH, Mogensen CE, Jensen HA, Evrin PE: Increased urinary albumin-excretion rate in benign essential hypertension. *Lancet* 1974, 1:1190-1192.
32. Knight EL, Kramer HM, Curhan GC: High-normal blood pressure and microalbuminuria. *Am.J.Kidney Dis.* 2003, 41:588-595.
33. Verdecchia P, Reboldi GP: Hypertension and microalbuminuria: the new detrimental duo. *Blood Press* 2004, 13:198-211.

34. Leoncini G, Sacchi G, Viazzi F, Ravera M, Parodi D, Ratto E, Vettoretti S, Tomolillo C, Deferrari G, Pontremoli R: Microalbuminuria identifies overall cardiovascular risk in essential hypertension: an artificial neural network-based approach. *J.Hypertens.* 2002, 20:1315-1321.
35. Bidani AK, Griffin KA: Pathophysiology of hypertensive renal damage: implications for therapy. *Hypertension* 2004, 44:595-601.
36. Grisk O, Rettig R: Interactions between the sympathetic nervous system and the kidneys in arterial hypertension. *Cardiovasc.Res.* 2004, 61:238-246.
37. Pedrinelli R, Penno G, Dell'Omo G, Bandinelli S, Giorgi D, Di B, V, Navalesi R, Mariani M: Microalbuminuria and transcapillary albumin leakage in essential hypertension. *Hypertension* 1999, 34:491-495.
38. Bohlen L, de Court, Weidmann P: Comparative study of the effect of ACE-inhibitors and other antihypertensive agents on proteinuria in diabetic patients. *Am.J.Hypertens.* 1994, 7:845-925.
39. Bakris GL, Fonseca V, Katholi RE, McGill JB, Messerli F, Phillips RA, Raskin P, Wright JT, Jr., Waterhouse B, Lukas MA, Anderson KM, Bell DS: Differential effects of beta-blockers on albuminuria in patients with type 2 diabetes. *Hypertension* 2005, 46:1309-1315.
40. Jerums G, Allen TJ, Campbell DJ, Cooper ME, Gilbert RE, Hammond JJ, O'Brien RC, Raffaele J, Tsalamandris C: Long-term renoprotection by perindopril or nifedipine in non-hypertensive patients with Type 2 diabetes and microalbuminuria. *Diabet.Med.* 2004, 21:1192-1199.
41. White WB, Duprez D, St Hillaire R, Krause S, Roniker B, Kuse-Hamilton J, Weber MA: Effects of the selective aldosterone blocker eplerenone versus the calcium antagonist amlodipine in systolic hypertension. *Hypertension* 2003, 41:1021-1026.
42. Brantsma AH, Bakker SJ, de Zeeuw D, de Jong PE, Gansevoort RT: Urinary albumin excretion as a predictor of the development of hypertension in the general population. *J.Am.Soc.Nephrol.* 2006, 17:331-335.
43. Mogensen CE: Prediction of clinical diabetic nephropathy in IDDM patients. Alternatives to microalbuminuria? *Diabetes* 1990, 39:761-767.
44. Schena FP, Gesualdo L: Pathogenetic mechanisms of diabetic nephropathy. *J.Am. Soc.Nephrol.* 2005, 16 Suppl 1:S30-S33.
45. Bohlender JM, Franke S, Stein G, Wolf G: Advanced glycation end products and the kidney. *Am.J.Physiol Renal Physiol* 2005, 289:F645-F659.
46. Ferrannini E, Buzzigoli G, Bonadonna R, Giorico MA, Oleggini M, Graziadei L, Pedrinelli R, Brandi L, Bevilacqua S: Insulin resistance in essential hypertension. *N.Engl.J.Med.* 1987, 317:350-357.
47. Bianchi S, Bigazzi R, Valtriani C, Chiapponi I, Sgherri G, Baldari G, Natali A, Ferrannini E, Campese VM: Elevated serum insulin levels in patients with essential hypertension and microalbuminuria. *Hypertension* 1994, 23:681-687.
48. Grandi AM, Santillo R, Zanzi P, Broggi R, Imperiale D, Colombo S, Bertolini A, Jessula A, Selva E, Guasti L, Venco A: Microalbuminuria in never-treated hypertensives: lack of relationship to hyperinsulinemia and genetic predisposition to hypertension. *Am.J.Hypertens.* 2000, 13:353-358.
49. Gall MA, Hougaard P, Borch-Johnsen K, Parving HH: Risk factors for development of incipient and overt diabetic nephropathy in patients with non-insulin dependent diabetes mellitus: prospective, observational study. *BMJ* 1997, 314:783-788.
50. Ravid M, Brosh D, Ravid-Safran D, Levy Z, Rachmani R: Main risk factors for nephropathy in type 2 diabetes mellitus are plasma cholesterol levels, mean blood pressure, and hyperglycemia. *Arch.Intern.Med.* 1998, 158:998-1004.

51. Chaturvedi N, Fuller JH, Taskinen MR: Differing associations of lipid and lipoprotein disturbances with the macrovascular and microvascular complications of type 1 diabetes. *Diabetes Care* 2001, 24:2071-2077.
52. Calvino J, Calvo C, Romero R, Gude F, Sanchez-Guisande D: Atherosclerosis profile and microalbuminuria in essential hypertension. *Am.J.Kidney Dis.* 1999, 34:996-1001.
53. Campese VM, Bianchi S, Bigazzi R: Association between hyperlipidemia and microalbuminuria in essential hypertension. *Kidney Int.Suppl* 1999, 71:S10-S13.
54. Cirillo M, Senigalliesi L, Laurenzi M, Alfieri R, Stamler J, Stamler R, Panarelli W, De Santo NG: Microalbuminuria in nondiabetic adults: relation of blood pressure, body mass index, plasma cholesterol levels, and smoking: The Gubbio Population Study. *Arch.Intern.Med.* 1998, 158:1933-1939.
55. Klahr S, Schreiner G, Ichikawa I: The progression of renal disease. *N.Engl.J.Med.* 1988, 318:1657-1666.
56. Tonolo G, Ciccarese M, Brizzi P, Puddu L, Secchi G, Calvia P, Atzeni MM, Melis MG, Maioli M: Reduction of albumin excretion rate in normotensive microalbuminuric type 2 diabetic patients during long-term simvastatin treatment. *Diabetes Care* 1997, 20:1891-1895.
57. Orth SR: Effects of smoking on systemic and intrarenal hemodynamics: influence on renal function. *J.Am.Soc.Nephrol.* 2004, 15 Suppl 1:S58-S63.
58. Halimi JM, Philippon C, Mimran A: Contrasting renal effects of nicotine in smokers and non-smokers. *Nephrol.Dial.Transplant.* 1998, 13:940-944.
59. Chase HP, Garg SK, Marshall G, Berg CL, Harris S, Jackson WE, Hamman RE: Cigarette smoking increases the risk of albuminuria among subjects with type I diabetes. *JAMA* 1991, 265:614-617.
60. Corradi L, Zoppi A, Tettamanti F, Malamani G, Lazzari P, Fogari R: Association between smoking and micro-albuminuria in hypertensive patients with type 2 diabetes mellitus. *J.Hypertens.Suppl* 1993, 11:S190-S191.
61. Gerstein HC, Mann JF, Pogue J, Dinneen SF, Halle JP, Hoogwerf B, Joyce C, Rashkow A, Young J, Zinman B, Yusuf S: Prevalence and determinants of microalbuminuria in high-risk diabetic and nondiabetic patients in the Heart Outcomes Prevention Evaluation Study. The HOPE Study Investigators. *Diabetes Care* 2000, 23 Suppl 2: B35-B39.
62. Pinto-Sietsma SJ, Mulder J, Janssen WM, Hillege HL, de Zeeuw D, de Jong PE: Smoking is related to albuminuria and abnormal renal function in nondiabetic persons. *Ann.Intern.Med.* 2000, 133:585-591.
63. Hillege HL, Janssen WM, Bak AA, Diercks GF, Grobbee DE, Crijns HJ, van Gilst WH, de Zeeuw D, de Jong PE: Microalbuminuria is common, also in a nondiabetic, nonhypertensive population, and an independent indicator of cardiovascular risk factors and cardiovascular morbidity. *J.Intern.Med.* 2001, 249:519-526.
64. Orth SR, Ritz E: The renal risks of smoking: an update. *Curr.Opin.Nephrol. Hypertens.* 2002, 11:483-488.
65. Cleland SJ, Reid JL: The renin-angiotensin system and the heart: a historical review. *Heart* 1996, 76:7-12.
66. Harris P: Evolution and the cardiac patient. *Cardiovasc.Res.* 1983, 17:437-445.
67. Skeggs LT, Lentz KE, Kahn JR, Shumway NP, Woods KR: The amino acid sequence of hypertensin. II. *J.Exp.Med.* 1956, 104:193-197.
68. Coates D: The angiotensin converting enzyme (ACE). *Int.J.Biochem.Cell Biol.* 2003, 35:769-773.

69. Struthers AD, MacDonald TM: Review of aldosterone- and angiotensin II-induced target organ damage and prevention. *Cardiovasc.Res.* 2004, 61:663-670.
70. de Gasparo M, Catt KJ, Inagami T, Wright JW, Unger T: International union of pharmacology. XXIII. The angiotensin II receptors. *Pharmacol.Rev.* 2000, 52:415-472.
71. Kambayashi Y, Bardhan S, Takahashi K, Tsuzuki S, Inui H, Hamakubo T, Inagami T: Molecular cloning of a novel angiotensin II receptor isoform involved in phosphotyrosine phosphatase inhibition. *J.Biol.Chem.* 1993, 268:24543-24546.
72. Murphy TJ, Alexander RW, Griendling KK, Runge MS, Bernstein KE: Isolation of a cDNA encoding the vascular type-1 angiotensin II receptor. *Nature* 1991, 351:233-236.
73. Campbell DJ: Tissue renin-angiotensin system: sites of angiotensin formation. *J.Cardiovasc.Pharmacol.* 1987, 10 Suppl 7:S1-S8.
74. Dzau VJ: Circulating versus local renin-angiotensin system in cardiovascular homeostasis. *Circulation* 1988, 77:14-13.
75. Bader M, Peters J, Baltatu O, Muller DN, Luft FC, Ganten D: Tissue renin-angiotensin systems: new insights from experimental animal models in hypertension research. *J.Mol.Med.* 2001, 79:76-102.
76. Zhuo JL, Imig JD, Hammond TG, Orengo S, Benes E, Navar LG: Ang II accumulation in rat renal endosomes during Ang II-induced hypertension: role of AT(1) receptor. *Hypertension* 2002, 39:116-121.
77. Ichikawa I, Harris RC: Angiotensin actions in the kidney: renewed insight into the old hormone. *Kidney Int.* 1991, 40:583-596.
78. Dzau VJ, Re R: Tissue angiotensin system in cardiovascular medicine. A paradigm shift? *Circulation* 1994, 89:493-498.
79. Wang T, Giebisch G: Effects of angiotensin II on electrolyte transport in the early and late distal tubule in rat kidney. *Am.J.Physiol* 1996, 271:F143-F149.
80. Navar LG, Harrison-Bernard LM, Imig JD, Wang CT, Cervenka L, Mitchell KD: Intrarenal angiotensin II generation and renal effects of AT1 receptor blockade. *J.Am.Soc.Nephrol.* 1999, 10 Suppl 12:S266-S272.
81. Wolf G: Angiotensin II as a mediator of tubulointerstitial injury. *Nephrol.Dial. Transplant.* 2000, 15 Suppl 6:61-63.
82. Ruiz-Ortega M, Ruperez M, Esteban V, Rodriguez-Vita J, Sanchez-Lopez E, Carvajal G, Egido J: Angiotensin II: a key factor in the inflammatory and fibrotic response in kidney diseases. *Nephrol.Dial.Transplant.* 2006, 21:16-20.
83. Hollenberg NK: Aldosterone in the development and progression of renal injury. *Kidney Int.* 2004, 66:1-9.
84. Bohrer MP, Deen WM, Robertson CR, Brenner BM: Mechanism of angiotensin II-induced proteinuria in the rat. *Am.J.Physiol* 1977, 233:F13-F21.
85. Lapinski R, Perico N, Remuzzi A, Sangalli F, Benigni A, Remuzzi G: Angiotensin II modulates glomerular capillary permselectivity in rat isolated perfused kidney. *J.Am.Soc.Nephrol.* 1996, 7:653-660.
86. Gomez-Garre D, Largo R, Tejera N, Fortes J, Manzarbeitia F, Egido J: Activation of NF-kappaB in tubular epithelial cells of rats with intense proteinuria: role of angiotensin II and endothelin-1. *Hypertension* 2001, 37:1171-1178.
87. Leehey DJ, Singh AK, Alavi N, Singh R: Role of angiotensin II in diabetic nephropathy. *Kidney Int.Suppl* 2000, 77:S93-S98.
88. Suzuki Y, Ruiz-Ortega M, Lorenzo O, Ruperez M, Esteban V, Egido J: Inflammation and angiotensin II. *Int.J.Biochem.Cell Biol.* 2003, 35:881-900.

89. Williams B, Baker AQ, Gallacher B, Lodwick D: Angiotensin II increases vascular permeability factor gene expression by human vascular smooth muscle cells. *Hypertension* 1995, 25:913-917.
90. Baldoncini R, Desideri G, Bellini C, Valenti M, De Mattia G, Santucci A, Ferri C: High plasma renin activity is combined with elevated urinary albumin excretion in essential hypertensive patients. *Kidney Int.* 1999, 56:1499-1504.
91. Pontremoli R, Sofia A, Tirota A, Ravera M, Nicoletta C, Viazzi F, Bezante GP, Borgia L, Bobola N, Ravazzolo R, Sacchi G, Deferrari G: The deletion polymorphism of the angiotensin I-converting enzyme gene is associated with target organ damage in essential hypertension. *J.Am.Soc.Nephrol.* 1996, 7:2550-2558.
92. Kario K, Kanai N, Nishiuma S, Fujii T, Saito K, Matsuo T, Matsuo M, Shimada K: Hypertensive nephropathy and the gene for angiotensin-converting enzyme. *Arterioscler.Thromb.Vasc.Biol.* 1997, 17:252-256.
93. Hadjadj S, Belloum R, Bouhanick B, Gallois Y, Guilloteau G, Chatellier G, Alhenc-Gelas F, Marre M: Prognostic value of angiotensin-I converting enzyme I/D polymorphism for nephropathy in type 1 diabetes mellitus: a prospective study. *J.Am.Soc.Nephrol.* 2001, 12:541-549.
94. Pontremoli R, Ravera M, Viazzi F, Nicoletta C, Berruti V, Leoncini G, Giacomelli F, Bezante GP, Sacchi G, Ravazzolo R, Deferrari G: Genetic polymorphism of the renin-angiotensin system and organ damage in essential hypertension. *Kidney Int.* 2000, 57:561-569.
95. Standards of Medical Care in Diabetes. *Diabetes Care* 2005, 28:S4-S36.
96. de Zeeuw D, Remuzzi G, Parving HH, Keane WF, Zhang Z, Shahinfar S, Snapinn S, Cooper ME, Mitch WE, Brenner BM: Proteinuria, a target for renoprotection in patients with type 2 diabetic nephropathy: lessons from RENAAL. *Kidney Int.* 2004, 65:2309-2320.
97. Engel SL, Schaeffer TR, Waugh MH, Rubin B: Effects of the nonapeptide SQ 20881 on blood pressure of rats with experimental renovascular hypertension. *Proc.Soc. Exp.Biol.Med.* 1973, 143:483-487.
98. Gavras H, Brunner HR, Laragh JH, Sealey JE, Gavras I, Vukovich RA: An angiotensin converting-enzyme inhibitor to identify and treat vasoconstrictor and volume factors in hypertensive patients. *N.Engl.J.Med.* 1974, 291:817-821.
99. Gansevoort RT, Sluiter WJ, Hemmelder MH, de Zeeuw D, de Jong PE: Antiproteinuric effect of blood-pressure-lowering agents: a meta-analysis of comparative trials. *Nephrol.Dial.Transplant.* 1995, 10:1963-1974.
100. Should all patients with type 1 diabetes mellitus and microalbuminuria receive angiotensin-converting enzyme inhibitors? A meta-analysis of individual patient data. *Ann.Intern.Med.* 2001, 134:370-379.
101. Lewis EJ, Hunsicker LG, Bain RP, Rohde RD: The effect of angiotensin-converting-enzyme inhibition on diabetic nephropathy. The Collaborative Study Group. *N.Engl.J.Med.* 1993, 329:1456-1462.
102. Viberti G, Mogensen CE, Groop LC, Pauls JF: Effect of captopril on progression to clinical proteinuria in patients with insulin-dependent diabetes mellitus and microalbuminuria. European Microalbuminuria Captopril Study Group. *JAMA* 1994, 271:275-279.
103. Effects of ramipril on cardiovascular and microvascular outcomes in people with diabetes mellitus: results of the HOPE study and MICRO-HOPE substudy. Heart Outcomes Prevention Evaluation Study Investigators. *Lancet* 2000, 355:253-259.

104. Ruggenenti P, Fassi A, Ilieva AP, Bruno S, Iliev IP, Brusegan V, Rubis N, Gherardi G, Arnoldi F, Ganeva M, Ene-Iordache B, Gaspari F, Perna A, Bossi A, Trevisan R, Dodesini AR, Remuzzi G: Preventing microalbuminuria in type 2 diabetes. *N.Engl. J.Med.* 2004, 351:1941-1951.
105. Dalla VM, Pozza G, Mosca A, Grazioli V, Lapolla A, Fioretto P, Crepaldi G: Effect of lercanidipine compared with ramipril on albumin excretion rate in hypertensive Type 2 diabetic patients with microalbuminuria: DIAL study (diabete, ipertensione, albuminuria, lercanidipina). *Diabetes Nutr.Metab* 2004, 17:259-266.
106. Randomised placebo-controlled trial of effect of ramipril on decline in glomerular filtration rate and risk of terminal renal failure in proteinuric, non-diabetic nephropathy. The GISEN Group (Gruppo Italiano di Studi Epidemiologici in Nefrologia). *Lancet* 1997, 349:1857-1863.
107. Ruggenenti P, Perna A, Benini R, Bertani T, Zoccali C, Maggiore Q, Salvadori M, Remuzzi G: In chronic nephropathies prolonged ACE inhibition can induce remission: dynamics of time-dependent changes in GFR. Investigators of the GISEN Group. Gruppo Italiano Studi Epidemiologici in Nefrologia. *J.Am.Soc.Nephrol.* 1999, 10:997-1006.
108. Staessen JA, Wang JG, Thijs L: Cardiovascular protection and blood pressure reduction: a meta-analysis. *Lancet* 2001, 358:1305-1315.
109. Fox KM: Efficacy of perindopril in reduction of cardiovascular events among patients with stable coronary artery disease: randomised, double-blind, placebo-controlled, multicentre trial (the EUROPA study). *Lancet* 2003, 362:782-788.
110. Wing LM, Reid CM, Ryan P, Beilin LJ, Brown MA, Jennings GL, Johnston CI, McNeil JJ, Macdonald GJ, Marley JE, Morgan TO, West MJ: A comparison of outcomes with angiotensin-converting-enzyme inhibitors and diuretics for hypertension in the elderly. *N.Engl.J.Med.* 2003, 348:583-592.
111. Nissen SE, Tuzcu EM, Libby P, Thompson PD, Ghali M, Garza D, Berman L, Shi H, Buebendorf E, Topol EJ: Effect of antihypertensive agents on cardiovascular events in patients with coronary disease and normal blood pressure: the CAMELOT study: a randomized controlled trial. *JAMA* 2004, 292:2217-2225.
112. Dahlof B, Sever PS, Poulter NR, Wedel H, Beevers DG, Caulfield M, Collins R, Kjeldsen SE, Kristinsson A, McInnes GT, Mehlsen J, Nieminen M, O'Brien E, Ostergren J: Prevention of cardiovascular events with an antihypertensive regimen of amlodipine adding perindopril as required versus atenolol adding bendroflumethiazide as required, in the Anglo-Scandinavian Cardiac Outcomes Trial-Blood Pressure Lowering Arm (ASCOT-BPLA): a multicentre randomised controlled trial. *Lancet* 2005, 366:895-906.
113. Pals DT, Masucci FD, Sipos F, Denning GS, Jr.: A specific competitive antagonist of the vascular action of angiotensin. II. *Circ.Res.* 1971, 29:664-672.
114. K/DOQI clinical practice guidelines on hypertension and antihypertensive agents in chronic kidney disease. *Am.J.Kidney Dis.* 2004, 43:S1-290.
115. Burnier M, Zanchi A: Blockade of the renin-angiotensin-aldosterone system: a key therapeutic strategy to reduce renal and cardiovascular events in patients with diabetes. *J.Hypertens.* 2006, 24:11-25.
116. Berl T: Angiotensin-converting enzyme inhibitors versus AT1 receptor antagonist in cardiovascular and renal protection: the case for AT1 receptor antagonist. *J.Am. Soc.Nephrol.* 2004, 15 Suppl 1:S71-S76.
117. Lansang MC, Stevanovic R, Price DA, Laffel LM, Hollenberg NK: ACE and non-ACE pathways in the renal vascular response to RAS interruption in type 1 diabetes mellitus. *Kidney Int.* 2005, 67:1033-1037.

118. Buter H, Navis G, Dullaart RP, de Zeeuw D, de Jong PE: Time course of the antiproteinuric and renal haemodynamic responses to losartan in microalbuminuric IDDM. *Nephrol.Dial.Transplant.* 2001, 16:771-775.
119. Parving HH, Lehnert H, Brochner-Mortensen J, Gomis R, Andersen S, Arner P: The effect of irbesartan on the development of diabetic nephropathy in patients with type 2 diabetes. *N.Engl.J.Med.* 2001, 345:870-878.
120. Lewis EJ, Hunsicker LG, Clarke WR, Berl T, Pohl MA, Lewis JB, Ritz E, Atkins RC, Rohde R, Raz I: Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N.Engl.J.Med.* 2001, 345:851-860.
121. Brenner BM, Cooper ME, de Zeeuw D, Keane WF, Mitch WE, Parving HH, Remuzzi G, Snapinn SM, Zhang Z, Shahinfar S: Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N.Engl.J.Med.* 2001, 345:861-869.
122. Viberti G, Wheeldon NM: Microalbuminuria reduction with valsartan in patients with type 2 diabetes mellitus: a blood pressure-independent effect. *Circulation* 2002, 106:672-678.
123. Barnett AH, Bain SC, Bouter P, Karlberg B, Madsbad S, Jervell J, Mustonen J: Angiotensin-receptor blockade versus converting-enzyme inhibition in type 2 diabetes and nephropathy. *N.Engl.J.Med.* 2004, 351:1952-1961.
124. Ziai F, Ots M, Provoost AP, Troy JL, Rennke HG, Brenner BM, Mackenzie HS: The angiotensin receptor antagonist, irbesartan, reduces renal injury in experimental chronic renal failure. *Kidney Int.Suppl* 1996, 57:S132-S136.
125. Nakaya H, Sasamura H, Kitamura Y, Amemiya T, Konishi K, Hayashi M, Saruta T: Effects of angiotensin inhibitors on renal injury and angiotensin receptor expression in early hypertensive nephrosclerosis. *Hypertens.Res.* 1999, 22:303-312.
126. Plum J, Buntin B, Nemeth R, Grabensee B: Effects of the angiotensin II antagonist valsartan on blood pressure, proteinuria, and renal hemodynamics in patients with chronic renal failure and hypertension. *J.Am.Soc.Nephrol.* 1998, 9:2223-2234.
127. Gansevoort RT, de Zeeuw D, de Jong PE: Is the antiproteinuric effect of ACE inhibition mediated by interference in the renin-angiotensin system? *Kidney Int.* 1994, 45:861-867.
128. Chan JC, Critchley JA, Tomlinson B, Chan TY, Cockram CS: Antihypertensive and anti-albuminuric effects of losartan potassium and felodipine in Chinese elderly hypertensive patients with or without non-insulin-dependent diabetes mellitus. *Am.J.Nephrol.* 1997, 17:72-80.
129. Nielsen S, Døllerup J, Nielsen B, Jensen HA, Mogensen CE: Losartan reduces albuminuria in patients with essential hypertension. An enalapril controlled 3 months study. *Nephrol.Dial.Transplant.* 1997, 12 Suppl 2:19-23.
130. Holdaas H, Hartmann A, Berg KJ, Lund K, Fauchald P: Renal effects of losartan and amlodipine in hypertensive patients with non-diabetic nephropathy. *Nephrol.Dial. Transplant.* 1998, 13:3096-3102.
131. Hermida RC, Calvo C, Ayala DE, Lopez JE: Decrease in urinary albumin excretion associated with the normalization of nocturnal blood pressure in hypertensive subjects. *Hypertension* 2005, 46:960-968.
132. Ishimitsu T, Kobayashi T, Honda T, Takahashi M, Minami J, Ohta S, Inada H, Yoshii M, Ono H, Matsuoka H: Protective effects of an angiotensin II receptor blocker and a long-acting calcium channel blocker against cardiovascular organ injuries in hypertensive patients. *Hypertens.Res.* 2005, 28:351-359.

133. Vogt L, Navis G, Koster J, Manolis AJ, Reid JL, de Zeeuw D: The angiotensin II receptor antagonist telmisartan reduces urinary albumin excretion in patients with isolated systolic hypertension: results of a randomized, double-blind, placebo-controlled trial. *J.Hypertens.* 2005, 23:2055-2061.
134. Ibsen H, Wachtell K, Olsen MH, Borch-Johnsen K, Lindholm LH, Mogensen CE, Dahlöf B, Devereux RB, de Faire U, Fyhrquist F, Julius S, Kjeldsen SE, Lederballe-Pedersen O, Nieminen MS, Omvik P, Oparil S, Wan Y: Does albuminuria predict cardiovascular outcome on treatment with losartan versus atenolol in hypertension with left ventricular hypertrophy? A LIFE substudy. *J.Hypertens.* 2004, 22:1805-1811.
135. Benson SC, Pershadsingh HA, Ho CI, Chittiboyina A, Desai P, Pravenec M, Qi N, Wang J, Avery MA, Kurtz TW: Identification of telmisartan as a unique angiotensin II receptor antagonist with selective PPARgamma-modulating activity. *Hypertension* 2004, 43:993-1002.
136. Fliser D, Buchholz K, Haller H: Antiinflammatory effects of angiotensin II subtype 1 receptor blockade in hypertensive patients with microinflammation. *Circulation* 2004, 110:1103-1107.
137. 2003 European Society of Hypertension-European Society of Cardiology guidelines for the management of arterial hypertension. *J.Hypertens.* 2003, 21:1011-1053.
138. Greene EL, Kren S, Hostetter TH: Role of aldosterone in the remnant kidney model in the rat. *J.Clin.Invest* 1996, 98:1063-1068.
139. Quan ZY, Walser M, Hill GS: Adrenalectomy ameliorates ablative nephropathy in the rat independently of corticosterone maintenance level. *Kidney Int.* 1992, 41:326-333.
140. Rocha R, Stier CT, Jr., Kifor I, Ochoa-Maya MR, Rennke HG, Williams GH, Adler GK: Aldosterone: a mediator of myocardial necrosis and renal arteriopathy. *Endocrinology* 2000, 141:3871-3878.
141. Horiuchi M, Nishiyama H, Hama J, Takenaka T, Kondo H, Kino H, Nagata S, Sugimura K, Katori R: Characterization of renal aldosterone receptors in genetically hypertensive rats. *Am.J.Physiol* 1993, 264:F286-F291.
142. Conn JW, Knoff RF, Nesbit RM: Clinical characteristics of primary aldosteronism from an analysis of 145 cases. *Am.J.Surg.* 1964, 107:159-172.
143. Chrysostomou A, Becker G: Spironolactone in addition to ACE inhibition to reduce proteinuria in patients with chronic renal disease. *N.Engl.J.Med.* 2001, 345:925-926.
144. Sato A, Hayashi K, Naruse M, Saruta T: Effectiveness of aldosterone blockade in patients with diabetic nephropathy. *Hypertension* 2003, 41:64-68.
145. Rossing K, Schjoedt KJ, Smidt UM, Boomsma F, Parving HH: Beneficial effects of adding spironolactone to recommended antihypertensive treatment in diabetic nephropathy: a randomized, double-masked, cross-over study. *Diabetes Care* 2005, 28:2106-2112.
146. Hayashi M, Tsutamoto T, Wada A, Tsutsui T, Ishii C, Ohno K, Fujii M, Taniguchi A, Hamatani T, Nozato Y, Kataoka K, Morigami N, Ohnishi M, Kinoshita M, Horie M: Immediate administration of mineralocorticoid receptor antagonist spironolactone prevents post-infarct left ventricular remodeling associated with suppression of a marker of myocardial collagen synthesis in patients with first anterior acute myocardial infarction. *Circulation* 2003, 107:2559-2565.
147. Pitt B, Zannad F, Remme WJ, Cody R, Castaigne A, Perez A, Palensky J, Wittes J: The effect of spironolactone on morbidity and mortality in patients with severe heart failure. Randomized Aldactone Evaluation Study Investigators. *N.Engl.J.Med.* 1999, 341:709-717.

148. Macdonald JE, Kennedy N, Struthers AD: Effects of spironolactone on endothelial function, vascular angiotensin converting enzyme activity, and other prognostic markers in patients with mild heart failure already taking optimal treatment. *Heart* 2004, 90:765-770.
149. Epstein M: Aldosterone receptor blockade and the role of eplerenone: evolving perspectives. *Nephrol.Dial.Transplant.* 2003, 18:1984-1992.
150. White WB, Duprez D, St Hillaire R, Krause S, Roniker B, Kuse-Hamilton J, Weber MA: Effects of the selective aldosterone blocker eplerenone versus the calcium antagonist amlodipine in systolic hypertension. *Hypertension* 2003, 41:1021-1026.
151. Brown NJ: Eplerenone: cardiovascular protection. *Circulation* 2003, 107:2512-2518.
152. Pitt B, Remme W, Zannad F, Neaton J, Martinez F, Roniker B, Bittman R, Hurley S, Kleiman J, Gattlin M: Eplerenone, a selective aldosterone blocker, in patients with left ventricular dysfunction after myocardial infarction. *N.Engl.J.Med.* 2003, 348:1309-1321.
153. Kobayashi N, Yoshida K, Nakano S, Ohno T, Honda T, Tsubokou Y, Matsuoka H: Cardioprotective mechanisms of eplerenone on cardiac performance and remodeling in failing rat hearts. *Hypertension* 2006, 47:671-679.
154. Wood JM, Maibaum J, Rahuel J, Grutter MG, Cohen NC, Rasetti V, Ruger H, Goschke R, Stutz S, Fuhrer W, Schilling W, Rigollier P, Yamaguchi Y, Cumin F, Baum HP, Schnell CR, Herold P, Mah R, Jensen C, O'Brien E, Stanton A, Bedigian MP: Structure-based design of aliskiren, a novel orally effective renin inhibitor. *Biochem.Biophys.Res.Comm.* 2003, 308:698-705.
155. Stanton A, Jensen C, Nussberger J, O'Brien E: Blood pressure lowering in essential hypertension with an oral renin inhibitor, aliskiren. *Hypertension* 2003, 42:1137-1143.
156. Gradman AH, Schmieder RE, Lins RL, Nussberger J, Chiang Y, Bedigian MP: Aliskiren, a novel orally effective renin inhibitor, provides dose-dependent antihypertensive efficacy and placebo-like tolerability in hypertensive patients. *Circulation* 2005, 111:1012-1018.
157. Nussberger J, Wuerzner G, Jensen C, Brunner HR: Angiotensin II suppression in humans by the orally active renin inhibitor Aliskiren (SPP100): comparison with enalapril. *Hypertension* 2002, 39:E1-E8.
158. Azizi M, Menard J, Bissery A, Guyenne TT, Bura-Riviere A, Vaidyanathan S, Camisasca RP: Pharmacologic demonstration of the synergistic effects of a combination of the renin inhibitor aliskiren and the AT1 receptor antagonist valsartan on the angiotensin II-renin feedback interruption. *J.Am.Soc.Nephrol.* 2004, 15:3126-3133.
159. Pilz B, Shagdarsuren E, Wellner M, Fiebeler A, Dechend R, Gratzke P, Meiners S, Feldman DL, Webb RL, Garrelds IM, Jan Danser AH, Luft FC, Muller DN: Aliskiren, a human renin inhibitor, ameliorates cardiac and renal damage in double-transgenic rats. *Hypertension* 2005, 46:569-576.
160. Fisher ND, Hollenberg NK: Renin inhibition: what are the therapeutic opportunities? *J.Am.Soc.Nephrol.* 2005, 16:592-599.

